
Superconductivity in two dimensions in the AlO_x/SrTiO₃ heterostructure

Shamashis Sengupta^{*1}, Miguel Monteverde², Tobias Rodel, Emilie Tisserond³, Florence Linez, Anil Murani⁴, Richard Deblock⁵, Sophie Gueron, Helene Bouchiat⁴, Philippe Lecoeur⁶, Thomas Maroutian, Claire Marrache-Kikuchi, Andrés Santander-Syro⁷, and Franck Fortuna

¹Centre de Sciences Nucléaires et de Sciences de la Matière (CSNSM) – CNRS : UMR8609, IN2P3, Université Paris XI - Paris Sud – Bâtiments 104 et 108 91405 Orsay Campus, France

²Laboratoire de Physique des Solides (LPS) – Université Paris-Sud – Bat 510, Centre Universitaire, France

³Laboratoire de Physique des Solides d'Orsay (LPS) – CNRS : UMR8502 – Université Paris-Sud Bâtiment 510 91405 ORSAY Cedex, France

⁴Laboratoire de Physique des Solides (LPS) – CNRS : UMR8502 – Université Paris-Sud, Laboratoire de Physique des Solides 91405 Orsay, France

⁵Laboratoire de Physique des Solides (LPS) – CNRS : UMR8502, Université Paris Sud - Paris XI – Université Paris-Sud, Laboratoire de Physique des Solides 91405 Orsay, France

^{6*} Institut d'Electronique Fondamentale, Université Paris-Sud, CNRS, UMR 8622, F-91405 Orsay, France, (IEF) – UMR 8622 – Bât. 220 - Centre Scientifique d'Orsay - F 91405 Orsay cedex, France

⁷Centre de Sciences Nucléaires et de Sciences de la Matière (CSNSM) – CNRS : UMR8609, IN2P3, Université Paris XI - Paris Sud – Université Paris-Sud Bâtiments 104 et 108 91405 Orsay Campus, France

Abstract

The realization of two-dimensional electronic gases (2DEGs) in oxide-based heterostructures has led to important discoveries [1,2] about superconductivity in low dimensions, such as the observation of pairing interactions without superconductivity [3] and density-of-states features resembling the pseudogap in cuprates [4]. Consequently, this 2DEG has emerged as a model system to study the physics of Cooper pair formation in two dimensions and to gain useful insights about complex problems, e.g., the phase diagram of high temperature superconductors.

In this talk, we will focus on: i) the facile realization of a 2DEG in AlO_x/SrTiO₃ using a method devised in our group that is more accessible to researchers than conventional hetero-epitaxial techniques, and ii) the phenomenon of gate-tunable superconductivity in two dimensions.

A conducting 2DEG can be realized on the bare surface of SrTiO₃ by the creation of oxygen vacancies [5]. The electrons populate the bands arising from Ti 3d orbitals. In a recent work (Rodel *et al.*, *Advanced Materials*, 28:1976–1980, 2016), we have shown that the deposition in ultra-high vacuum of a thin layer of metallic Al on SrTiO₃ leads to the creation of the

*Speaker

same 2DEG due to the withdrawal of oxygen atoms from the surface by the reducing agent Al (which turns into insulating AlOx). The resulting heterostructure, AlOx/SrTiO3, is suitable for transport experiments because the layer of AlOx protects the 2DEG in ambient conditions against the percolation of oxygen from air. From transport experiments, we have determined that the 2DEG is superconducting with a critical temperature of 320 mK. The critical parameters (temperature and field) are tunable with the gate voltage, leading to a 'superconducting dome' in the phase diagram. Phase fluctuations of the order parameter play a dominant role in the phenomenon of superconductivity in two dimensions. By continuously varying the carrier density, we have been able to tune the 2DEG across different regimes of electronic interactions that influence the phenomenon of Cooper pair condensation. We will present our results on this topic.

References:

Reyren et al., Science 317, 1196 (2007)

Cavaglia et al., Nature 456, 624 (2008)

Cheng et al., Nature 521, 196 (2015)

Richter et al., Nature 502, 528 (2013)

Santander-Syro et al., Nature 469, 189 (2011)